

## Homework #6. Due Thursday, October 18th, in class

### Reading:

1. For this homework assignment. In Pugh: 2.5; in Rudin: 2.5 and 4.4; class notes (Lecture 12).
2. For next week's classes: Continuity and connectedness (2.5 in Pugh and 4.4 in Rudin). Completions of metric spaces (2.10 in Pugh). We will also start talking about uniform convergence (4.1 in Pugh and 7.1-7.3 in Rudin).

### Problems:

**Note on hints:** All hints are given at the end of the assignment, each on a separate page. Problems (or parts of problems) for which hint is available are marked with \*.

1. Complete the proof of the backwards direction of Theorem 12.2 from class (which asserts the any interval in  $\mathbb{R}$  is connected). Recall that we
  - (i) proved the result for closed bounded intervals (intervals of the form  $[a, b]$ ) and
  - (ii) explained how to prove the result for intervals of the form  $[a, b)$  using (i) and Lemma 12.5 (=Problem 4 in this assignment).

Prove the backwards direction of Theorem 12.2 in the remaining cases.

**2\*.** Let  $X$  be a metric space. Prove that  $X$  is disconnected if and only if there exists a continuous function  $f : X \rightarrow \mathbb{R}$  such that  $f(X) = \{1, -1\}$ .

**3.**

(a\*) Let  $X$  be a disconnected metric space, so that  $X = A \sqcup B$  for some non-empty closed subsets  $A$  and  $B$ . Prove that if  $C$  is any connected subset of  $X$ , then  $C \subseteq A$  or  $C \subseteq B$ .

(b\*) A metric space  $X$  is called *path-connected* if for any  $x, y \in X$  there exists a continuous function  $f : [0, 1] \rightarrow X$  such that  $f(0) = x$  and  $f(1) = y$  (informally, this means that any two points in  $X$  can be joined by a path in  $X$ ). Prove that any path-connected metric space is connected.

**4\*.** Let  $X$  be a metric space,  $\{X_\alpha\}_{\alpha \in I}$  a collection (not necessarily finite) of subsets of  $X$  such that  $\bigcap_{\alpha \in I} X_\alpha$  is non-empty and  $\bigcup_{\alpha \in I} X_\alpha = X$ . Prove that if each  $X_\alpha$  is connected, then  $X$  is connected.

**5.** (practice) Metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  are called **isometric** if there exists a bijection  $f : X \rightarrow Y$  such that  $d_Y(f(a), f(b)) = d_X(a, b)$  for

all  $a, b \in X$ . Prove that all abstract properties of metric spaces introduced in this class are preserved under isometries, that is, if  $(X, d_X)$  and  $(Y, d_Y)$  are isometric and  $X$  is compact, then  $Y$  is compact; if  $X$  is connected, then  $Y$  is connected etc.

**6.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces, and consider the product space  $X \times Y$  with metric  $d$  given by  $d((x_1, y_1), (x_2, y_2)) = d_X(x_1, x_2) + d_Y(y_1, y_2)$

- (a) Prove that  $(X \times Y, d)$  is indeed a metric space
- (b) Prove that for every  $x \in X$ , the subset  $\{x\} \times Y = \{(x, y) : y \in Y\}$  of  $X \times Y$  is isometric to  $Y$ . Likewise for every  $y \in Y$ , the subset  $X \times \{y\} = \{(x, y) : x \in X\}$  is isometric to  $X$ .
- (c\*) Prove that if  $X$  and  $Y$  are both connected, then  $X \times Y$  connected.

**7.** The goal of this problem is to prove that any open subset of  $\mathbb{R}$  (with standard metric) is a **disjoint** union of at most countably many open intervals.

So, let  $U$  be any open subset of  $\mathbb{R}$ .

- (a) Define the relation  $\sim$  on  $U$  by setting  $x \sim y \iff x = y$  or  $(x < y$  and  $[x, y] \subset U)$  or  $(y < x$  and  $[y, x] \subset U)$ . Prove that  $\sim$  is an equivalence relation.
- (b\*) Let  $A$  be an equivalence class with respect to  $\sim$ . Show that  $A$  is an open interval.
- (c\*) Deduce from (b) that  $U$  is a disjoint union of open intervals. Then prove that the number of those intervals is at most countable.

**8.** Use Problem 6 to show that the analogue of Problem 7 does not hold in  $\mathbb{R}^2$ , that is, there exist open subsets of  $\mathbb{R}^2$  which are not representable as disjoint unions of open discs (an open disc is an open ball in  $\mathbb{R}^2$ ).

**Hint for 2:** The “ $\Leftarrow$ ” direction is easy. For the “ $\Rightarrow$ ” direction, assume that  $X = A \sqcup B$  with  $A, B$  closed, and show that the function  $f : X \rightarrow \mathbb{R}$  given by

$$f(x) = \begin{cases} 1 & \text{if } x \in A \\ -1 & \text{if } x \in B \end{cases}$$

is continuous.

4

**Hint for 3(a):** Use the inheritance principle.

**Hint for 3(b):** Use (a) and Theorem 12.1 from class (=Theorem 4.22 from Rudin).

**Hint for 4:** Assume that  $X$  is disconnected and use 3(a) to reach a contradiction.

**Hint for 6(b):** By 5 and 6(a) all subsets of the form  $X \times \{y\}$  and  $\{x\} \times Y$  are connected. Start with this fact and use Problem 4 twice. Drawing a picture in the case  $X = Y = [0, 1]$  will likely be helpful.

**Hint for 7(b):** First use an earlier homework problem to prove that  $A$  is connected. Then use Theorem 12.2 from class.

**Hint for 7(c):** Use the fact that  $\mathbb{Q}$  is countable.