

Homework #7. Due Thursday, October 30th, in class

Reading:

1. For this homework assignment: 4.5-4.7, 7.2-7.5 + class notes (Lectures 12-16)
2. For next week's classes: Section 7.6 (equicontinuous families of functions). I plan to follow Rudin quite closely in this section.

Note: In some problems you may want to use the following stronger version of Theorem 14.1 from class: *Let X be a metric space, and fix $a \in X$. Let $\{f_n\}_{n \in \mathbb{N}}$, f be functions from X to \mathbb{R} , and suppose that $f_n \rightrightarrows f$ and each f_n is continuous at a . Then f is also continuous at a .*

(In class we showed that global continuity of each f_n implies global continuity of f , but we did not explicitly state that continuity of each f_n at a single point a forces f to be continuous at the same point; however, the proof given in class actually establishes the above stronger form of Theorem 14.1).

Problems:

1. Consider functions $f_n : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ given by $f_n(x) = \frac{1}{nx+1}$. Let $0 \leq a \leq b$ be real numbers. Prove that $\{f_n\}$ converges uniformly on $[a, b] \iff a > 0$ or $a = b = 0$.

2. Let X be a set and $\{f_n\}, f$ functions from X to \mathbb{R} . Suppose that $f_n \rightrightarrows f$ on X , f is bounded on X (that is, there exists $M \in \mathbb{R}$ such that $|f(x)| \leq M$ for all $x \in X$) and also that each f_n is bounded. Prove that the sequence $\{f_n\}$ is uniformly bounded (see Def. 7.19 in Rudin). **Hint:** it may be useful to draw a picture.

3. Let X be a metric space and $\{f_n\}, f$ functions from X to \mathbb{R} . Suppose that $f_n \rightrightarrows f$ on X and each f_n is uniformly continuous. Prove that f is uniformly continuous. **Hint:** Imitate the proof of Theorem 14.1.

4. For each $\alpha \in \mathbb{R}$ define the function $I_\alpha : \mathbb{R} \rightarrow \mathbb{R}$ by

$$I_\alpha(x) = \begin{cases} 0 & \text{if } x < \alpha \\ 1 & \text{if } x \geq \alpha \end{cases}$$

Now let $S = \{s_1, s_2, \dots\}$ be a countable infinite subset of \mathbb{R} , and define $f : \mathbb{R} \rightarrow \mathbb{R}$ by $f(x) = \sum_{n=1}^{\infty} \frac{I_{s_n}(x)}{2^n}$. Prove that this series always converges (so that f is indeed defined on \mathbb{R}), f is increasing, and f is continuous at $x \iff x \notin S$. **Hint:** Use Weierstrass M-test.

5. Let $f : [a, b] \rightarrow \mathbb{R}$ be an increasing function. Prove that the following are equivalent:

- (i) f is continuous (on $[a, b]$)
- (ii) $f([a, b]) = [f(a), f(b)]$
- (iii) $f([a, b])$ is dense in $[f(a), f(b)]$

Hint: It is easy to show that (i) \Rightarrow (ii) \Rightarrow (iii). To prove that (iii) \Rightarrow (i) use the results about discontinuities of increasing functions proved in class.

6. Give a short proof of Theorem 7.17 from Rudin under the additional assumption that each f'_n is continuous (naturally, you should not be repeating or imitating the proof in the general case given in Rudin). **Hint:** in the notations of Theorem 7.17, if we set $g(x) = \lim_{n \rightarrow \infty} f'_n(x)$, then g is a uniform limit of continuous functions, hence g is also continuous (Theorem 14.1). Now define $f : [a, b] \rightarrow \mathbb{R}$ by $f(x) = \int_{x_0}^x g(t) dt + C$ for suitable C and prove that $f_n \rightrightarrows f$ using the Fundamental Theorem of Calculus (recall that $\int_c^d g(t) dt$ in the case $c > d$ is defined as $-\int_d^c g(t) dt$).

7. Let X be a any set.

- (a) Let $B(X)$ be the set of all bounded functions $f : X \rightarrow \mathbb{R}$. Define the map $d : B(X) \times B(X) \rightarrow \mathbb{R}_{\geq 0}$ by

$$d(f, g) = \sup_{x \in X} |f(x) - g(x)|.$$

Prove that d is a metric on $B(X)$ which satisfies the following property: given $f_n, f \in B(X)$, we have $f_n \rightrightarrows f$ (on X) $\iff d(f_n, f) \rightarrow 0$ as $n \rightarrow \infty$. Equivalently, $f_n \rightrightarrows f$ (on X) $\iff f_n \rightarrow f$ in the metric space $(B(X), d)$. (This was stated but not proved at the end of Lecture 15).

- (b) (optional) The goal of this part is to establish the analog of (a) with $B(X)$ replaced by $Func(X, \mathbb{R})$, the set of all functions from $X \rightarrow \mathbb{R}$.

Let $\Omega = Func(X, \mathbb{R})$, and define $D : \Omega \times \Omega \rightarrow \mathbb{R}_{\geq 0}$ by

$$D(f, g) = \min\{\sup_{x \in X} |f(x) - g(x)|, 1\}.$$

Prove that D is a metric on Ω such that given $\{f_n\}, f \in \Omega$, we have $f_n \rightrightarrows f$ on X $\iff f_n \rightarrow f$ in the metric space (Ω, D) .